



8-epi-PGF2 α ELISA Kit

12/19

(Catalog # E4789-100; 96 assays, Storage at 4°C)

I. Introduction:

F2-isoprostanes are a family of prostaglandin F2-like compounds that are formed by free-radical-catalyzed peroxidation of arachidonic acid. Several F2-isoprostanes, but in particular 8-epi PGF2 α , are widely used as oxidative stress biomarkers. High levels of F2-IPs can disrupt the physicochemical integrity of cell membranes but more importantly can exert significant pharmacological activity. 8-epi-PGF2 α is potent vasoconstrictor acting on the renal, pulmonary, and retinal vasculatures, modulates platelet aggregation and is involved in the activation of intracellular signaling. BioVision's 8-epi-PGF2 α ELISA Kit is based on the Competitive ELISA principle. The micro-plate provided in this kit has been pre-coated with 8-epi-PGF2 α . During the reaction, 8-epi-PGF2 α in the samples or standard competes with 8-epi-PGF2 α coated on the plate for binding to the anti-8-epi-PGF2 α antibody. Then Horseradish Peroxidase (HRP) conjugate is added to each microplate well, and TMB substrate is for color development. There is a negative correlation between the OD value of samples and the concentration of 8-epi-PGF2 α . The concentration of 8-epi-PGF2 α in the samples can be calculated by comparing the OD of the samples to the standard curve.

II. Applications:

in vitro quantitative measurement of 8-epi-PGF2 α in serum, plasma and other biological samples.

Sensitivity: 9.38 pg/mL

Detection Range: 15.63—1000 pg/mL

Specificity: This kit recognizes 8-epi-PGF2 α in samples. No Significant cross-reactivity or interference between 8-epiPGF2 α and analogues was observed.

Precision: Coefficient of variation is < 10%.

III. Sample Type:

Serum, plasma and other biological fluids

IV. Kit Contents:

Components	E4789-100	Part Number	Storage
Micro ELISA Plate	8 wells x12 strips	E4789-100-1	-20°C
Reference Standard	2 vials	E4789-100-2	-20°C
Biotinylated Detection Ab (100x)	120 μ l	E4789-100-3	-20°C
HRP Conjugate (100x)	120 μ l	E4789-100-4	-20°C (protect from light)
Reference Standard & Sample Diluent	20 ml	E4789-100-5	4°C
Biotinylated Detection Antibody Diluent	14 ml	E4789-100-6	4°C
HRP Conjugate Diluent	14 ml	E4789-100-7	4°C
Wash Buffer (25X)	30 ml	E4789-100-8	4°C
Substrate Reagent	10 ml	E4789-100-9	4°C (protect from light)
Stop Solution	10 ml	E4789-100-10	4°C
Plate Sealer	4	E4789-100-11	4°C

V. User Supplied Reagents and Equipment:

- Microplate reader capable of measuring absorbance at 450 nm
- Clean Eppendorf tubes for preparing standards or sample dilutions

VI. Storage and Handling:

Store at 4°C.

VII. Reagent and Sample Preparation:

Bring all reagents to room temperature before use. Before using the kit, spin tubes and bring down all components to the bottom of tubes.

- **Wash Buffer (25X):** Dilute 30 ml of Concentrated Wash Buffer with 720 ml of deionized or distilled water to prepare 750 ml of Wash Buffer. (Note: if crystals have formed in the concentrate, warm it in a 40°C water bath and mix it gently until the crystals have completely dissolved)
- **Biotinylated Detection Antibody working solution:** Calculate the required amount (50 μ L/well). Centrifuge the stock tube before use, dilute the 100x Concentrated Biotinylated Detection Antibody to 1x working solution with Biotinylated Detection Antibody Diluent.
- **HRP Conjugate working solution:** Calculate the required amount before the experiment (100 μ L/well). In preparation, slightly more than calculated should be prepared. Dilute the 100x Concentrated HRP Conjugate to 1x working solution with Concentrated HRP Conjugate Diluent.
- **Standard:** Centrifuge the standard at 10,000xg for 1 min. Add 1.0 ml of Standard and Sample Diluent, let it stand for 10 min and invert it gently several times. After it dissolves fully, mix it thoroughly with a pipette. This reconstitution produces a working solution of 1000

