



EZClick™ Palmitoylated Protein Assay Kit (FACS/Microscopy), Green Fluorescence

rev 07/20

(Catalog # K452-100; 100 assays; Store at -20°C)

I. Introduction:

Palmitoylation occurs when fatty acids, such as Palmitic Acid are covalently attached to the side chains of cysteine (and less frequently to serine, threonine side chains) in proteins. This type of post-translational modification greatly affects cellular trafficking, compartmentalization and membrane tethering. Palmitoyltransferase (PAT), the enzyme responsible for this modification transfers a palmitate moiety from palmitoyl-CoA to the thiol group of cysteine in the target protein. Compared to myristoylation, palmitoylation is reversible and the reverse reaction is catalyzed by thioesterases. The Palmitoylation/Depalmitoylation cycle plays an important role when modified protein shuttles between cellular compartments. PAT mutations are associated with many neurological diseases and cancer progression. **BioVision offers EZClick™ Palmitoylated Protein Assay Kit** is a highly specific, simple and robust method for labeling and detection of palmitoylated proteins. The kit uses a modified Palmitic Acid that is fed directly into the cells and gets incorporated into proteins during or post translation. This post translational modification can be followed by click reaction with an azide-containing dye. The assay kit offers a powerful method for imaging localization, trafficking, and dynamics of Palmitoylated proteins or detection by FACS for quantitative studies. We provide sufficient materials for 100 assays in a 96-well plate format.

II. Applications:

- Identification and localization of Palmitic acid modified proteins
- Detection and quantification of biosynthesis, dynamics and turnover of palmitoylated proteins
- Screening for genotoxic compounds and effectors of protein modifications

III. Sample Type:

- Suspension or adherent cell cultures

IV. Kit Contents:

Components	K452-100	Cap Code	Part Number
EZClick™ Wash Buffer (10X)	25 ml	NM	K452-100-1
Fixative Solution	10 ml	WM	K452-100-2
Permeabilization Buffer (10X)	25 ml	Blue NM	K452-100-3
EZClick™ Palmitic Acid Label (1000X)	10 µl	White	K452-100-4
Copper Reagent (100X)	100 µl	Blue	K452-100-5
EZClick™ Fluorescent Azide (100X)	100 µl	Green/Amber	K452-100-6
Reducing Agent (20X)	500 µl	Yellow	K452-100-7
EZClick™ Total DNA Stain (1000X)	20 µl	Blue/Amber	K452-100-8

V. User Supplied Reagents and Equipment:

- Tissue culture vessels and appropriate culturing media
- Phosphate Buffered Saline (PBS, pH 7.4)
- Sterile 0.1% Gelatin Solution (optional, only required for suspension cells)
- Flow cytometer equipped with laser capable of excitation at 488 nm wavelength (FL-1)
- Fluorescence microscope capable of excitation and emission at 440/490 nm and UV filter

VI. Storage Conditions and Reagent Preparation:

Upon arrival, store the entire kit at -20°C protected from light. Briefly centrifuge small vials prior to opening. Read the entire protocol before performing the assay.

- **10X Wash Buffer and 10X Permeabilization Buffer: Thaw at 37°C to dissolve completely.** Dilute the 10X stocks at 1:10 in sterile water, mix well. Store at 4°C.
- **Fixative Solution: Divide into aliquots and store at -20°C, protected from light.**
- **Remaining components:** Store at -20°C protected from light. While in use, keep on ice and minimize light exposure.

VII. Palmitoylated Protein Assay Protocol:

Notes:

This assay was developed using HeLa (adherent) and Jurkat (suspension) cells and can be modified for any cell lines. The protocol below refers to a 96-well tissue culture plate. Adjust volumes accordingly for other plate formats. The assay volume is 100 µl. Growth conditions, cell numbers per well and other factors may affect the incorporation rate of the protein label. Therefore, optimize the assay for your cell type. We suggest an initial test of several EZClick™ Palmitic Acid Label concentrations to find the best condition for your experimental design. Avoid stressing the cells by washes or temperature changes prior to incubation with EZClick™ Palmitic Acid Label. All steps should be carried out at room temperature (RT) unless otherwise specified. Equilibrate all buffers to RT prior to the experiment.

1. Labeling of control and experimental cells: Method with drug pre-incubation:

- Obtain cell suspension of desired density and seed directly into tissue culture vessels, or on coverslips for high resolution microscopy. **To immobilize suspension cells for microscopy:** Add 100 µl of 0.1% gelatin solution directly into the wells. Tilt the plate to cover the entire well surface and place it in a tissue culture hood for 1 hr. Gently remove the gelatin solution and seed your cells. Allow the cells to recover overnight before the treatment. Next day, treat the cells with appropriate effectors according to your protocol. Do not add treatment to the positive and negative control cells. **Negative Control Cells** (Unstained Cells, cells not exposed to Palmitic Acid Label or EZClick™ Fluorescent Azide), **Background Control Cells** (Cells are not exposed to the EZClick™

