# BioVision

# **TRAF3 Polyclonal Antibody**

| CATALOG NO:         | 3563-30T<br>3563-100  | 30 µg (Trial size)<br>100 µg |
|---------------------|---|------------------------------|
| HOST:               | Rabbit  |                              |
| IMMUNOGEN:          | Synthetic peptide surrounding amino acid 326 of rat TRAF3 (ID# BV-F62G) |                              |
| SPECIES REACTIVITY: | Human, Mouse, Rat, Pig, House, Sheep, Cattle, Monkey                    |                              |

#### FORMULATION:

0.5 mg/ml affinity purified rabbit polyclonal antibody in phosphate-buffered saline (PBS) containing 30% glycerol, 0.5% BSA, 5 mM EDTA and 0.01% thimerosal.

#### STORAGE CONDITIONS:

Store at -20°C. For long-term storage, aliquot and freeze at -70°C. Avoid repeated freeze/defrost cycles.

#### BACKGROUND:


TRAFs (TNF receptor associated proteins) form a family of cytoplasmic adapter proteins that mediate signal transduction from many members of the TNF-receptor superfamily and the interleukin-1 receptor. The carboxy-terminal region of TRAFs is required for self-association and interaction with receptor cytoplasmic domains following ligand-induced oligomerization. Recent molecular cloning studies have lead to identification of six TRAFs (TRAF1-TRAF6). TRAF3, originally named CRAF1, interacts directly with the CD40 cytoplasmic tail through a region of similarity to the tumor necrosis factor-alpha (TNF-alpha) receptor-associated factors. TRAF3 binds only a single site, which contains the sequence PEQET, whereas TRAF1 and TRAF2 are capable of binding to either the PEQET site or an additional downstream domain.

# SPECIFICITY:

The antibody recognizes 67 kDa TRAF3 of human, mouse and rat origins. Reactivity to other species has not been tested.

# APPLICATION AND USAGE:

The antibody can be used in Western blotting (0.5-4 µg/ml). However, the optimal conditions should be determined individually. Blocking peptide (Cat.# 3563BP-50) is available separately.



Cat. #:3563-100 Western blot analysis of TRAF3 in lysates from Jurkat cells (Lane 1 & 2), 3T3 cells (Lane 3) and rat kidney (Lane 4).

### **RELATED PRODUCTS:**

Apoptosis Detection Kits & Reagents

- Annexin V Kits & Bulk Reagents
- Caspase Assay Kits & Reagents
- Mitochondrial Apoptosis Kits & Reagents
- Nuclear Apoptosis Kits & Reagents
- Apoptosis Inducers and Set
- Apoptosis siRNA Vectors

Cell Fractionation System

- Mitochondria/Cytosol Fractionation Kit
- Nuclear/Cytosol Fractionation Kit
- Membrane Protein Extraction Kit
- Cytosol/Particulate Rapid Separation Kit
- Mammalian Cell Extraction Kit
- FractionPREP Fractionation System

Cell Proliferation & Senescence

- Quick Cell Proliferation Assay Kit
- Senescence Detection Kit
- High Throughput Apoptosis/Cell Viability Assay Kits
- LDH-Cytotoxicity Assay Kit
- Bioluminescence Cytotoxicity Assay Kit
- Live/Dead Cell Staining Kit

Cell Damage & Repair

- HDAC Fluorometric & Colorimetric Assays & Drug Discovery Kits
- HAT Colorimetric Assay Kit & Reagents
- DNA Damage Quantification Kit
- Glutathione & Nitric Oxide Fluorometric & Colorimetric Assay Kits

Signal Transduction

- cAMP & cGMP Assay Kits
- Akt & JNK Activity Assay Kits
- Beta-Secretase Activity Assay Kit

Adipocyte & Lipid Transfer

- Recombinant Adiponectin, Survivin, & Leptin
- CETP Activity Assay & Drug Discovery Kits
- PLTP Activity Assay & Drug Discovery Kits
- Total Cholesterol Quantification Kit
- Molecular Biology & Reporter Assays
  - siRNA Vectors
  - Cloning Insert Quick Screening Kit
  - Mitochondrial & Genomic DNA Isolation Kits
  - 5 Minutes DNA Ligation Kit
  - 20 Minutes Gel Staining/Destaining Kit
  - β-Galactosidase Staining Kit & Luciferase Reporter Assay Kit

Growth Factors and Cytokines

Monoclonal and Polyclonal Antibodies

