

H4pan Antibody

ALTERNATE NAMES: Histone H4

CATALOG #: 6825-25

AMOUNT: 25 µg

HOST/ISOTYPE: Rabbit

IMMUNOGEN: KLH-conjugated synthetic peptide of Histone H4 containing an unmodified sequence from the central part of the protein.

FORM: Liquid

FORMULATION: In PBS with 0.05% (W/V) sodium azide and 0.05% ProClin 300.

PURIFICATION: Affinity purified.

SPECIES REACTIVITY: Human.


STORAGE CONDITIONS: Store at -20°C; for long storage, store at -80°C. Avoid multiple freeze-thaw cycles.

DESCRIPTION: Histones are the main constituents of the protein part of chromosomes of eukaryotic cells. They are rich in the amino acids arginine and lysine and have been greatly conserved during evolution. Histones pack the DNA into tight masses of chromatin. Two core histones of each class H2A, H2B, H3 and H4 assemble and are wrapped by 146 base pairs of DNA to form one octameric nucleosome. Histones play a central role in the regulation of transcription, DNA repair, DNA replication and chromosomal stability. These different functions are established via a complex set of post-translational modifications which either directly or indirectly alter chromatin structure and DNA accessibility to facilitate transcriptional activation or repression or other nuclear processes.


APPLICATION: ELISA: 1:500, Dot Blot: 1:10,000, ChIP: 2-5 µg/ChIP.

Note: This information is only intended as a guide. The optimal dilutions must be determined by the user.

08/14



ChIP assays were performed using HeLa cells and the antibody and optimized PCR primer sets for qPCR. A titration of the antibody consisting of 1, 2, 5, and 10 µl per ChIP experiment was analysed. IgG (5 µg/IP) was used as negative control. The Fig shows the recovery, expressed as a % of input (the relative amount of IP DNA compared to input DNA after qPCR analysis).




RELATED PRODUCTS:

- H3R17me2 Antibody (Cat # 6803-50)
- H3K9me1 Antibody (Cat # 6804-50)
- H3K36me2 Antibody (Cat # 6805-50)
- H3 Pan Antibody (Cat # 6806-50)
- H4K8ac Antibody (Cat # 6807-50)

FOR RESEARCH USE ONLY! Not to be used on humans.

For research use only



To determine the titer, an ELISA was performed using a serial dilution of the antibody. The antigen used was a peptide containing the histone modification of interest. By plotting the absorbance against the antibody dilution the titer of the antibody was estimated to be 1:11250.

A Dot Blot analysis was performed to test the cross reactivity of the antibody with the peptide used for immunization of the rabbit and other peptides containing unmodified sequences of different histones. 100 to 0.2 pmol of the peptide containing the respective histone modification were spotted on a membrane. The Fig shows a high specificity of the antibody for the modification of interest.

