

## Phospho-IGF1R (Tyr980) Polyclonal Antibody

<b>CATALOG NO:</b>	A1040-100
<b>ALTERNATIVE NAMES:</b>	IGF-I receptor, CD221, IGF1R
<b>AMOUNT:</b>	100 µg (1 mg/ml)
<b>HOST/ISOTYPE:</b>	Rabbit IgG
<b>IMMUNOGEN:</b>	Synthetic peptide
<b>SPECIES REACTIVITY:</b>	Human
<b>PURIFICATION:</b>	Affinity purified rabbit IgG using protein G
<b>FORM:</b>	Liquid
<b>FORMULATION:</b>	Supplied in 0.01M Tris-HCl, pH 8.0, 0.15M NaCl, and 0.02% sodium azide. Protein was determined by OD280 absorbance.
<b>STORAGE CONDITIONS:</b>	Store at 4°C. For long term storage, aliquot and freeze at -20°C. Avoid repeated freeze/thaw cycles.
<b>DESCRIPTION:</b>	Receptor tyrosine kinase which mediates actions of insulin-like growth factor 1 (IGF1). Binds IGF1 with high affinity and IGF2 and insulin (INS) with a lower affinity. The activated IGF1R is involved in cell growth and survival control. IGF1R is crucial for tumor transformation and survival of malignant cell. Ligand binding activates the receptor kinase, leading to receptor autophosphorylation, and tyrosines phosphorylation of multiple substrates, that function as signaling adapter proteins including, the insulin-receptor substrates (IRS1/2), Shc and 14-3-3 proteins. Phosphorylation of IRSs proteins lead to the activation of two main signaling pathways: the PI3K-AKT/PKB pathway and the Ras-MAPK pathway. The result of activating the MAPK pathway is increased cellular proliferation, whereas activating the PI3K pathway inhibits apoptosis and stimulates protein synthesis. Phosphorylated IRS1 can activate the 85 kDa regulatory subunit of PI3K (PIK3R1), leading to activation of several downstream substrates, including protein AKT/PKB. AKT phosphorylation, in turn, enhances protein synthesis through mTOR activation and triggers the antiapoptotic effects of IGF1R through phosphorylation and inactivation of BAD. In parallel to PI3K-driven signaling, recruitment of Grb2/SOS by phosphorylated IRS1 or Shc leads to recruitment of Ras and activation of the ras-MAPK pathway. In addition to these two main signaling pathways IGF1R signals also through the Janus kinase/signal transducer and activator of transcription pathway (JAK/STAT). Phosphorylation of JAK proteins can lead to phosphorylation/activation of signal transducers and activators of transcription (STAT) proteins. In particular activation of STAT3, may be essential for the transforming activity of IGF1R. The JAK/STAT pathway activates gene transcription and may be responsible for the transforming

activity. JNK kinases can also be activated by the IGF1R. IGF1 exerts inhibiting activities on JNK activation via phosphorylation and inhibition of MAP3K5/ASK1, which is able to directly associate with the IGF1R.

**APPLICATION:** Western blot  
Immunoblot: 0.5-2 µg/ml  
ELISA

**Note: This information is only intended as a guide. The optimal dilutions must be determined by the user.**

### RELATED PRODUCTS:

- Human CellExp™ IGF1R/CD221, human recombinant (**Cat. No. 7490-10**)
- IGF-1 sR (human) ELISA Kit (**Cat. No. K4776-100**)
- IGF-1 (human) ELISA Kit (**Cat. No. K4775-100**)
- IGF-1, human recombinant (**Cat. No. 4119**)
- IGF-1, murine recombinant (**Cat. No. 4120**)
- IGF-1, rat recombinant (**Cat. No. 4121**)
- R3 IGF-1, human recombinant (**Cat. No. 4216**)
- Human CellExp™ IGF-1, human recombinant (**Cat. No. 7507**)
- Anti-Rat IGF-1 Antibody (**Cat. No. 5121-100**)
- IGF-1 Antibody (**Cat. No. 5119-100**)
- IGF-1 Antibody (**Cat. No. 5120R-100**)
- IGF-I (human) ELISA Kit (**Cat. No. K4775**)
- IGF-1, human recombinant (**Cat. No. 4119-100**)
- Anti-Rat IGF-1 Antibody (**Cat. No. 5121-100**)
- R3 IGF-1, human recombinant (**Cat. No. 4216-1G**)

**FOR RESEARCH USE ONLY! Not to be used on humans.**