

Anti-HIST1H1E K16ac Antibody

CATALOG NO.: A2056-100 (100 µl)

BACKGROUND DESCRIPTION: Histones are basic nuclear proteins responsible for nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H1 family. Transcripts from this gene lack poly A tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6.

ALTERNATE NAMES: Histone H1.4 (Histone H1b) (Histone H1s-4), HIST1H1E, H1F4

ANTIBODY TYPE: Polyclonal

HOST/ISOTYPE: Rabbit / IgG

IMMUNOGEN: Acetylated peptide sequence targeting residues around Lysine 16 of human Histone H1.4

PURIFICATION: Antigen Affinity purified

FORM: Liquid


FORMULATION: In 0.01 M PBS, 50% glycerol, 0.03% proclin 300, pH 7.4

SPECIES REACTIVITY: Human


STORAGE CONDITIONS: Store at -20°C. Avoid freeze / thaw cycles

APPLICATIONS AND USAGE: ICC 1:20-1:200, IF 1:50-1:200


Note: This information is only intended as a guide. The optimal dilutions must be determined by the user



Immunofluorescence analysis of MCF-7 cells (treated with sodium butyrate, 30 mM, 4 hrs) using Anti-HIST1H1E K16ac antibody at dilution of 1:100 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG (H+L) as secondary antibody.



Immunocytochemistry analysis of MCF-7 cells using Anti-HIST1H1E K16ac antibody at dilution of 1:100.



ChIP analysis of HeLa cells (4×10^6 , treated with 30 mM sodium butyrate for 4 hrs) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with 8 μ g Anti-HIST1H1E K16ac antibody or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the β -Globin promoter.

RELATED PRODUCTS:

H3K27ac polyclonal antibody (6869)

H3K9ac polyclonal antibody (6872)

Histone H4 Antibody (3624)

H4K5ac polyclonal antibody (6820)

FOR RESEARCH USE ONLY! Not to be used on humans.