

## **Emetine dihydrochloride hydrate**

ALTERNATE NAME: (2S,3R,11bS)-2-[[(1R)-6,7-dimethoxy-1,2,3,4-tetrahydroisoquinolin-1-yl]methyl]-3-ethyl-9,10-dimethoxy-

2,3,4,6,7,11b-hexahydro-1H-benzo[a]quinolizine, dihydrochloride, hydrate

Emetine hydrochloride hydrate

**CATALOG #**: B2339-250

B2339-50

**AMOUNT:** 250 mg 50 mg

STRUCTURE:

H,CO NH OCH,

H—CI NH<sub>2</sub>O NH<sub>3</sub>O

**MOLECULAR FORMULA:**  $C_{29}H_{40}N_2O_4.2HCl. xH_2O$ 

MOLECULAR WEIGHT: 553.56 (anhydrous basis)

**CAS NUMBER:** 7083-71-8

APPEARANCE: White to off-white solid

PURITY: ≥98% by HPLC

SOLUBILITY: >100 mg/ml (H<sub>2</sub> O)

STORAGE: 4°C

REFERENCES: Visnyei, K., et al. (2011). Mol. Cancer Ther. 10, 1818-1828.

**DESCRIPTION:** An alkaloid derived from ipecac root. Irreversibly blocks protein synthesis by inhibiting the movement of

ribosome along the mRNA. Induces hypotension by blocking adrenoreceptors. Inhibits DNA replication in the early S phase. Inhibits HIF-1 activation by hypoxia. Induces apoptosis in leukemia cells. Also, selectively inhibits multiple glioblastoma (GBM) stem cells-enriched cultures. It has been found to inhibit

endemic human coronaviruses and MERS-CoV in cell culture and cell entry assays.

HANDLING: Do not take internally. Wear gloves and mask when handling the product! Avoid contact by all modes of

exposure.

**RELATED PRODUCTS:** 

Clindamycin hydrochloride (2627)
EZSolution Puromycin Dihydrochloride (1861)
Emetine dihydrochloride (1970)
6-Azauridine (B1024)
Streptomycin Sulfate (2517)
Cycloheximide (1041)
Puromycin dihydrochloride (1860)
G418 Sulfate (1557)
Kanamycin Sulfate (2498)
Anisomycin (1549)

Blasticidin S Hydrochloride (1859)

DISCLAIMER: FOR RESEARCH USE ONLY! Not to be used on humans.