

Bovine Serum Albumin – Heat Shock, Protease DNASE Free, pH 7.0

CATALOG #:	7920-5	5 g
	7920-25	25 g
	7920-100	100 g
	7920-1000	1 kg

SOURCE: The raw material used to manufacture this lot was derived exclusively from USDA-regulated abattoirs located in the United States. The animals received ante-mortem and post-mortem inspection under a licensed veterinarian's supervision and were certified at the time of slaughter as free of infectious or contagious diseases and injurious parasites.

PURITY: ≥ 96% by SDS-PAGE analysis

MOL. WEIGHT: ~ 66.43 kDa

FORM: Off-white powder

OTHER PARAMETERS:

Albumin Content: ≥ 98% by electrophoresis.

pH: 7.0 ± 0.3

Moisture: ≤ 5.0%

Heavy Metals: ≤ 20 ppm

Ash: ≤ 2%

Protease: Not Detected

DNASE: Not Detected

IgG: Not Detected

SOLUBILITY: ~ 50 mg/ml in water, with 15 min manual stirring.

STORAGE CONDITIONS: This BSA is stable for ≥ 5 years when stored at +5°C +/- 3°C in a sealed dry container.

DESCRIPTION: This BSA was purified by a proprietary heat shock process to inactivate proteases. BSA makes up approximately 60% of all proteins in animal serum. It is commonly used in cell culture protocols, particularly when protein supplementation is necessary and the other components of serum are unwanted. It aids in stabilization of enzymes and other proteins to prevent proteolysis and denaturation, especially when the

proteins are present in low concentrations. In cell culture its main role is as a carrier of small molecules. Because of its negative charge, BSA binds water, salts, fatty acids, vitamins and hormones, then carries these bound components between tissues and cells. The binding capacity also makes BSA an effective scavenger for removing toxic substances, including pyrogens, from the medium.

APPLICATIONS/ADVANTAGES:

- Excellent in RIA and ELISA and as a diluent or blocking agent.
- No detectable protease and DNASE activity.

RELATED PRODUCTS:

- BSA – Cohn Fraction V, pH – 5.2 (**7904-5, 25, 100**)
- BSA – Cohn Fraction V, pH – 7.0 (**7905-5, 25, 100**)
- BSA – Cohn Fraction V, Immunoassay Grade, Protease Free, pH – 7.0 (**7906-5, 25, 100**)
- BSA – Cohn Fraction V, Fatty Acid Free, pH – 7.0 (**7907-5, 25, 100**)
- BSA – Cohn Fraction V, Fatty Acid Free, pH – 5.2 (**7908-5, 25, 100**)
- BSA – Cohn Fraction V, Endotoxin Low, pH – 7.0 (**7909-5, 25, 100**)
- BSA – Cohn Fraction V, Endotoxin Low, pH – 5.2 (**7910-5, 25, 100**)
- BSA – Cohn Fraction V, Immunoassay Grade, Protease Free, pH – 5.2 (**7911-5, 25, 100**)
- BSA – Cohn Fraction V, Fatty Acid Low, pH – 7.0 (**7912-5, 25, 100**)
- BSA (10% in H₂O) (**2119-10**)
- BSA Antibody (**5998-100**)
- AGE-BSA (**2221-10**)
- Glucose AGE-BSA (**2223-10**)
- Biotinylated BSA (Biotin-LC-BSA) (3 biotin/BSA) (**7097-5, 25**)
- Biotinylated BSA (Biotin-LC-BSA) (5 biotin/BSA) (**7098-5, 25**)
- Biotinylated BSA (Biotin-LC-BSA) (12 biotin/BSA) (**7099-5, 25**)
- Bovine Serum Albumin – Fraction V, 30% Solution (**7914-100, 500**)
- BSA – Heat Shock, pH – 7.0 (**7915-5, 25, 100**)
- BSA – Heat Shock, pH – 5.2 (**7916-5, 25, 100**)
- BSA – Heat Shock, Reagent Grade pH – 7.0 (**7917-5, 25, 100**)
- BSA – Heat Shock, Diagnostic Grade pH – 7.0 (**7918-5, 25, 100**)
- BSA – Heat Shock, Protease Free pH – 7.0 (**7919-5, 25, 100**)
- BSA – Heat Shock, Fatty Acid Free pH – 7.0 (**7921-5, 25, 100**)
- BSA – Heat Shock, Low Endotoxin pH – 7.0 (**7922-5, 25, 100**)

FOR RESEARCH USE ONLY! Not to be used in humans.