# **Recombinant Human BMPR1A**

**CATALOG #:** 4881-10 10 μg 4881-1000 1 mg

LOT #: \_\_\_\_\_

SYNONYMS: BMPR-1A, BMP-R1A, BMPR1A, BMR1A, CD292, CD-292

SOURCE: Insect cells (Baculovirus)

**PURITY:** > 90 % by SDS-PAGE and RP-HPLC analyses

**FORM:** Lyophilized from sterile PBS.

# **RECONSTITUTION:**

Centrifuge the vial prior to opening. Reconstitute in sterile PBS to a concentration  $\geq$  100 µg/ml. This solution can then be diluted into other aqueous buffers.

## STORAGE CONDITIONS:

The lyophilized protein is best-stored desiccated at -20°C. Reconstituted human BMPR1A should be stored at 4°C for 2-7 days and at -20°C for future use. For long term storage it is recommended to add a carrier protein (0.1 % HSA or BSA) and store as working aliquots at -20°C.

#### DESCRIPTION:

The bone morphogenetic protein (BMP) receptors are a family of transmembrane serine/threonine kinases that include the type I receptors BMPR1A and BMPR1B and the type II receptor BMPR2. These receptors are also closely related to the Activin receptors, ACVR1 and ACVR2. The ligands of these receptors are members of the TGF-beta superfamily. TGF-betas and activins transduce their signals through the formation of heteromeric complexes with 2 different types of serine (threonine) kinase receptors: type I receptors of about 50-55 kDa and type II receptors of about 70-80 kDa. Type II receptors bind ligands in the absence of type I receptors, but they require their respective type I receptors for signaling, whereas type I receptors require their respective type II receptors for ligand binding. BMPR1A Human Recombinant extracellular domain produced in baculovirus is a monomeric, glycosylated, polypeptide chain fused with 6xHis tag at C-terminus and having a molecular mass of 23 kDa. The BMR1A is purified by proprietary chromatographic techniques.

#### **BIOLOGICAL ACTIVTY:**

Measured by its ability to inhibit recombinant human BMP-2 induced alkaline phosphatase production by C2C12 myogenic cells. The ED $_{50}$  for this effect is typically 1-3  $\mu$ g/ml in the presence of 500 ng/ml of recombinant human BMP-2.

FOR RESEARCH USE ONLY! Not to be used in humans.

## **RELATED PRODUCTS:**

#### Apoptosis Detection Kits & Reagents

- Annexin V Kits & Bulk Reagents
- Caspase Assay Kits & Reagents
- Mitochondrial Apoptosis Kits & Reagents
- Nuclear Apoptosis Kits & Reagents
- Apoptosis Inducers and Set
- Apoptosis siRNA Vectors

## Cell Fractionation System

- Mitochondria/Cytosol Fractionation Kit
- Nuclear/Cytosol Fractionation Kit
- Membrane Protein Extraction Kit
- Cytosol/Particulate Rapid Separation Kit
- Mammalian Cell Extraction Kit
- FractionPREP Fractionation System

## Cell Proliferation & Senescence

- Quick Cell Proliferation Assay Kit
- Senescence Detection Kit
- High Throughput Apoptosis/Cell Viability Assay Kits
- LDH-Cytotoxicity Assay Kit
- Bioluminescence Cytotoxicity Assay Kit
- Live/Dead Cell Staining Kit

## Cell Damage & Repair

- HDAC Fluorometric & Colorimetric Assays & Drug Discovery Kits
- HAT Colorimetric Assay Kit & Reagents
- DNA Damage Quantification Kit
- Glutathione & Nitric Oxide Fluorometric & Colorimetric Assay Kits

# Signal Transduction

- cAMP & cGMP Assay Kits
- Akt & JNK Activity Assay Kits
- Beta-Secretase Activity Assay Kit

## Adipocyte & Lipid Transfer

- Recombinant Adiponectin, Survivin, & Leptin
- CETP Activity Assay & Drug Discovery Kits
- PLTP Activity Assay & Drug Discovery Kits
- Total Cholesterol Quantification Kit

## Molecular Biology & Reporter Assays

- siRNA Vectors
- Cloning Insert Quick Screening Kit
- Mitochondrial & Genomic DNA Isolation Kits
- 5 Minutes DNA Ligation Kit
- 20 Minutes Gel Staining/Destaining Kit
- β-Galactosidase Staining Kit & Luciferase Reporter Assay Kit

#### **Growth Factors and Cytokines**

Managland and Dalidand Anthodia



For research use only