

Recombinant Human IL-31

CATALOG #:	4983-10 4983-1000	10 µg 1 mg
LOT #:	_____	
SYNONYMS:	Interleukin 31, IL31, IL-31	
SOURCE:	<i>E. coli</i>	
PURITY:	> 95 % by SDS-PAGE and SEC-HPLC	
FORM:	Lyophilized from 20 mM phosphate buffer, 150 mM NaCl, pH 7.4	

RECONSTITUTION:

Centrifuge the vial prior to opening. Reconstitute in sterile ddH₂O to a concentration ≥ 100 µg/ml. This solution can then be diluted into other aqueous buffers.

STORAGE CONDITIONS:

The lyophilized protein is best-stored desiccated at -20°C. Reconstituted human IL-31 should be stored at 4°C for 2-7 days and at -20°C for future use. Avoid freeze/thaw cycles.

DESCRIPTION:

IL-31 produced by activated Th2-type T cells, cooperates with a heterodimeric receptor consisting of IL-31 Receptor Antagonist and Oncostatin-M Receptor that is constitutively expressed on epithelial cells and keratinocytes. IL-31 plays a role in the promotion of allergic skin disorders and in regulating other allergic diseases, such as asthma. IL-31 is involved in the itching sensation and endorses the scratching behavior in NC/Nga mice with atopic dermatitis. IL-31 expression is connected with CLA(+) T cells and contributes to the development of atopic dermatitis-induced skin inflammation and pruritus. IL-31 is a powerful inducer of proinflammatory mediators in human colonic SEMFs. IL-31 takes part as a proinflammatory cytokine derived from Th2 cells. Serum IL-31 level is higher in patients with atopic dermatitis. IL-31 is involved in a broad range of immune- & non-immune cells & possesses potential pleiotropic physiological functions, including regulating hematopoiesis & immune response, causing inflammatory bowel disease, airway hypersensitivity & dermatitis. IL-31 human recombinant produced in *E. coli* is a single, non-glycosylated, polypeptide chain containing 141 amino acids (24-164 a.a.) and having a molecular mass of 15.8 kDa.

BIOLOGICAL ACTIVITY:

The ED₅₀ was determined by its ability to activate STAT following receptor ligand interaction and found to be <5 ng/ml, corresponding to a specific activity of 200,000 units.

AMINO ACID SEQUENCE:

SHTLPVRLLRPSDDVQKIVEELQSLSKMLLKDVVEEEKGVLVSQNYTLPCLSFDAQPPNNIHSPA
IRAYLKTIKRLQDNKSVIDEIIEHLDKLIFQDAPETNISVPTDTHECKRFILTSQQFSECMDLALK
SLTSGAQQATT

FOR RESEARCH USE ONLY! Not to be used in humans.

RELATED PRODUCTS:

Apoptosis Detection Kits & Reagents

- Annexin V Kits & Bulk Reagents
- Caspase Assay Kits & Reagents
- Mitochondrial Apoptosis Kits & Reagents
- Nuclear Apoptosis Kits & Reagents
- Apoptosis Inducers and Set
- Apoptosis siRNA Vectors

Cell Fractionation System

- Mitochondria/Cytosol Fractionation Kit
- Nuclear/Cytosol Fractionation Kit
- Membrane Protein Extraction Kit
- Cytosol/Particulate Rapid Separation Kit
- Mammalian Cell Extraction Kit
- FractionPREP Fractionation System

Cell Proliferation & Senescence

- Quick Cell Proliferation Assay Kit
- Senescence Detection Kit
- High Throughput Apoptosis/Cell Viability Assay Kits
- LDH-Cytotoxicity Assay Kit
- Bioluminescence Cytotoxicity Assay Kit
- Live/Dead Cell Staining Kit

Cell Damage & Repair

- HDAC Fluorometric & Colorimetric Assays & Drug Discovery Kits
- HAT Colorimetric Assay Kit & Reagents
- DNA Damage Quantification Kit
- Glutathione & Nitric Oxide Fluorometric & Colorimetric Assay Kits

Signal Transduction

- cAMP & cGMP Assay Kits
- Akt & JNK Activity Assay Kits
- Beta-Secretase Activity Assay Kit

Adipocyte & Lipid Transfer

- Recombinant Adiponectin, Survivin, & Leptin
- CETP Activity Assay & Drug Discovery Kits
- PLTP Activity Assay & Drug Discovery Kits
- Total Cholesterol Quantification Kit

Molecular Biology & Reporter Assays

- siRNA Vectors
- Cloning Insert Quick Screening Kit
- Mitochondrial & Genomic DNA Isolation Kits
- 5 Minutes DNA Ligation Kit
- 20 Minutes Gel Staining/Destaining Kit
- β-Galactosidase Staining Kit & Luciferase Reporter Assay Kit

Growth Factors and Cytokines

Monoclonal and Polyclonal Antibodies