

## IL-17A, Human CellExp™, human recombinant

|                            |   |
|----------------------------|---|
| <b>CATALOG #:</b>          | 6468-10            10 µg<br>6468-50            50 µg  |
| <b>ALTERNATE NAMES:</b>    | CTLA-8, IL-17, IL-17A, Cytotoxic T-lymphocyte-associated antigen 8.                         |
| <b>SOURCE:</b>             | Human 293 Cell Expressed  |
| <b>PURITY:</b>             | > 95% by SDS - PAGE   |
| <b>MOL. WEIGHT:</b>        | 30 to 35 kDa, homodimer, glycosylated   |
| <b>ENDOTOXIN LEVEL:</b>    | < 1.0 EU per 1 µg of protein  |
| <b>FORMULATION:</b>        | Lyophilized in 10 mM Tris, pH 7.4 and 250 mM NaCl.  |
| <b>RECONSTITUTION:</b>     | Reconstitute in sterile PBS containing 0.1% endotoxin-free recombinant human serum albumin. |
| <b>STORAGE CONDITIONS:</b> | Aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.            |

**ADVANTAGES:**


- Animal-derived product free
- High Activity
- Authentic Glycosylation

**DESCRIPTION:**

Interleukin-17 is a proinflammatory cytokine produced by activated T cells. IL-17 regulates the activities of NF-kappa B and mitogen-activated protein kinases. IL17 can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of IL-17 are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. The original described IL-17 protein, now known as IL-17A, was identified from a CD4+ T cells DNA library. IL-17 can be induced from primary peripheral blood CD4+ T cells upon stimulation. Supernatant from cells transfected with IL-17 induced IL-16 and IL-18 production and enhanced the surface expression of the intracellular adhesion molecule-1 (ICAM-1) in human fibroblasts.

**BIOLOGICAL ACTIVITY:**

ED<sub>50</sub> is typically 5 to 25 ng/mL. The specific activity was determined by the dose-dependent induction of IL-6 secretion from NDHF Adult fibroblasts.



Human Cell<sup>exp</sup> Human Recombinant IL-17A

**RELATED PRODUCTS:**

- Human Cell<sup>exp</sup> Human Recombinant IL-17F (Cat # 6469-10, -50)
- IL-17, murine recombinant (Cat # 4177-10, -1000)
- IL-17A, human recombinant (Cat # 4176-25, -1000)
- IL-17A, rat recombinant (Cat # 4178-25, -1000)
- IL-17A/F, human recombinant (Cat # 4176AF-10, -50, -1000)
- IL-17E, human recombinant (Cat # 4176E-25, -1000)
- IL-17E, murine recombinant (Cat # 4177E-25, -100, -1000)
- IL-17E, rat recombinant (Cat # 4178E-25, -100, -1000)
- IL-17F, human recombinant (Cat # 4176F-25, -1000)
- IL-17F, murine recombinant (Cat # 4177F-25, -100, -1000)
- IL-17F, rat recombinant (Cat # 4178F-25, -100, -1000)
- IL-17 Antibody (Cat # 5176-200)

**FOR RESEARCH USE ONLY! Not to be used in humans.**

