BioVision 8/13 For research use only

sFas Receptor, Human Recombinant

CATALOG #: 7141-10 10 μg

7141-50 50 μg

ALTERNATE NAMES: soluble Fas receptor (sFasR), TNFRSF6, CD95,

Apo I, Fas Antigen

SOURCE: E Coli

PURITY: ≥ 98% by SDS-PAGE gel and HPLC analyses

MOL. WEIGHT: 17.6 kDa

ENDOTOXIN LEVEL: $< 0.1 \text{ ng/}\mu\text{g} \text{ of protein } (<1\text{EU/}\mu\text{g}).$

FORM: Lyophilized

FORMULATION: Sterile filtered through a 0.2 micron filter.

Lyophilized with no additives.

STORAGE CONDITIONS: Store at -20°C. After reconstitution, aliquot and

store at -20°C to -80°C. Avoid repeated freezing

and thawing cycles.

RECONSTITUTION:

Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 week. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

DESCRIPTION:

Fas and Fas Ligand (FasL) belong to the TNF superfamily and are type I and type II transmembrane proteins, respectively. Binding of FasL to Fas triggers apoptosis in Fasbearing cells. The mechanism of apoptosis involves recruitment of pro-caspase 8 through an adaptor molecule called FADD followed by processing of the pro-enzyme to active forms. These active caspases then cleave various cellular substrates leading to the eventual cell death. sFasR is capable of inhibiting FasL induced apoptosis by acting as a

decoy receptor that serves as a sink for FasL. The full length Fas (receptor) is a 319 amino acid type I transmembrane protein, which contains a 157 amino acid extracellular domain, a 17 amino acid transmembrane domain, and 145 amino acid cytoplasmic domain. Recombinant human soluble Fas (sFas Receptor) is a 157 amino acid polypeptide (17.6 kDa) corresponding to the TNFR homologous cysteine rich extracellular domain Fas.

BIOLOGICAL ACTIVITY: ED $_{50}$ was determined by its ability to inhibit the cytotoxicity of Jurkat cells using a concentration range of 10-15 μ g/ml in the presence of 2ng/ml of hFasL.

AMINO ACID SEQUENCE:

MRLSSKSVNA QVTDINSKGL ELRKTVTTVE TQNLEGLHHD GQFCHKPCPP GERKARDCTV NGDEPDCVPC QEGKEYTDKA HFSSKCRRCR LCDEGHGLEV EINCTRTQNT KCRCKPNFFC NSTVCEHCDP CTKCEHGIIK ECTLTSNTKC KEEGSRS

RELATED PRODUCTS:

- sFas Ligand, human recombinant (Cat. No. 7140-10, -50)
- FAS Inhibitor, C75 (Cat. No. 1547-1)
- Fas Ligand Antibody (Clone I-6060) (Cat. No. 3330-100)
- Fas/Apo1 Antibody (Cat. No. 3070R-100)
- Fas/Apo1 Blocking Peptide (Cat. No. 3070RBP-50)
- FasL Antibody (Cat. No. 3345R-100)
- FasL Blocking Peptide (Cat. No. 3345RBP-50)

FOR RESEARCH USE ONLY! Not to be used in humans.

