

Neuropoietin, murine recombinant

CATALOG #:	7180-10	10 µg
	7180-50	50 µg
ALTERNATE NAMES:	NPO, NP	
SOURCE:	E. Coli	
PURITY:	≥ 98% by SDS-PAGE gel and HPLC analyses	
MOL. WEIGHT:	19.8 kDa	
ENDOTOXIN LEVEL:	< 0.1 ng/µg of protein (<1EU/µg).	
FORM:	Lyophilized	
FORMULATION:	Sterile filtered through a 0.2 micron filter. Lyophilized from 2.5 mM Tris, pH 10.2 and 0.5 mM DTT	
STORAGE CONDITIONS:	Store at -20°C. After reconstitution, aliquot and store at -20°C to -80°C. Avoid repeated freezing and thawing cycles.	

RECONSTITUTION:

Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 week. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

DESCRIPTION:

Neuropoietin is a newly identified member of the IL-6 cytokine family. Members of this family, including IL-6, IL-11, oncostatin M, leukemia inhibitory factor (LIF), cardiotrophin-1 (CT-1), cardiotrophin-like cytokine, and CNTF, display a four-helix bundle structure, and signal through gp130-containing receptor complexes. Neuropoietin, which is predominantly expressed in neuroepithelia during embryonic life, acts through a receptor complex comprising CNTF receptor- α component, gp130, and LIF receptor. Like CNTF, it promotes the survival of embryonic motor neurons and could increase the proliferation of neural precursor cells in the presence of EGF and FGF-2. Interestingly, the human Neuropoietin gene has evolved toward a pseudogene, suggesting that the alternative signaling via CNTF is an effective compensatory pathway. Recombinant murine

AMINO ACID SEQUENCE:

MAPISPSEPI GQAYSLALYM QKNTSALLQT YLQHQQSPFS DPGFSAPELQ
LSTLPSAAVS FKTWHAMEDA ERLSRAQGAF LALTQHLQLV GDDQSYLNPG
SPILLAQLGA ARLRAQGLLG NMAAIMTALG LPIPEEDTL GFVPPFGASAF
ERKCRGYIVT REYGHWTDR VLDLALLKAK YSA

BIOLOGICAL ACTIVITY:

Determined by the dose dependent stimulation of the proliferation of human TF-1 cells. The expected ED₅₀ is 0.5-0.8 µg/ml.

RELATED PRODUCTS:

- IL-6, human recombinant (**Cat # 4143-20, -50, -1000**)
- IL-6, murine recombinant (**Cat # 4144-10, -50, -1000**)
- IL-6, rat recombinant (**Cat # 4145-10, -100, -1000**)
- IL-6 Antibody (**Cat # 5143-200**)
- IL-6 Antibody (**Cat # 5144-200**)
- IL-6 Antibody (**Cat # 5145-200**)

FOR RESEARCH USE ONLY! Not to be used in humans.