**BioVision** rev 02/14 For research use only

## PCSK9, Human CellExp™, murine recombinant

**CATALOG #:** 7266-20 20 μg

ALTERNATE NAMES: PCSK9, FH3, HCHOLA3, LDLCQ1, NARC-1,

NARC1, PC9, Proprotein convertase

subtilisin/kexin type 9

SOURCE: HEK 293 cells

**PURITY:** ≥ 97% by SDS-PAGE gel

MOL. WEIGHT: This protein is fused with 6×His tag at the N-

terminus, has a calculated MW of 72 kDa. The predicted N-terminus is Gln 35. DTT-reduced Protein migrates as 20 kDa and 64 kDa due to

glycosylation.

**ENDOTOXIN LEVEL:** < 0.1 ng/µg of protein (<1EU/µg) by LAL method

FORM: Lyophilized

**FORMULATION:** Sterile filtered through a 0.22 micron filter. Lyophilized from 1 x PBS, pH 7.4. Generally 5-8% Mannitol or trehalose is added as a protectant before lyophilization.

**STORAGE CONDITIONS:** Store at -20°C. After reconstitution, aliquot and store at -20°C and use within 3 months. Avoid repeated freezing and thawing cycles.


**RECONSTITUTION:** Centrifuge the vial prior to opening. Reconstitute in sterile PBS, pH 7.4 to a concentration of 100 μg/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 month. For extended storage, it is recommended to store at -20°C.

**DESCRIPTION**: Proprotein convertase subtilisin/kexin type 9 (PCSK9), is an enzyme which in humans is encoded by the PCSK9 gene. This gene encodes a proprotein convertase belonging to the proteinase K subfamily of the secretory subtilase family. This protein plays a major regulatory role in cholesterol homeostasis. PCSK9 binds to the epidermal growth factor-like repeat A (EGF-A) domain of the low-density lipoprotein receptor (LDLR), inducing LDLR degradation. PCSK9 may also have a role in the

differentiation of cortical neurons. Mutations in this gene have been associated with a rare form of autosomal dominant familial hypercholesterolemia (HCHOLA3).

## **BIOLOGICAL ACTIVITY:**

Measured by its binding ability in a functional ELISA Immobilized mouse PCSK9 at 10  $\mu$ g/ml (100  $\mu$ l/well) can bind biotinylated rhLDLR. The EC<sub>50</sub> of biotinylated human LDLR is 0.1  $\mu$ g/ml.



Murine recombinant PCSK9

## **RELATED PRODUCTS:**

- PCSK9, human recombinant (Cat # 7265-20)
- PCSK9 Antibody (Cat # 5112-100)

FOR RESEARCH USE ONLY! Not to be used in humans.

