**BioVision** 2/14 For research use only

## Cytidine deaminase, human recombinant

**CATALOG #:** 7363-100 100 μg

**ALTERNATE NAMES:** CDA, CDD, Cytidine aminohydrolase, Cytosine

nucleoside deaminase

SOURCE: E. coli

**PURITY:** > 90% by SDS-PAGE

FORM: Liquid

FORMULATION: 0.5 mg/ml in 20 mM Tris-HCl buffer (pH 8.0)

containing 100 mM NaCl, 1 mM DTT, 2 mM

EDTA and 40% glycerol.

**MOL. WEIGHT:** 18.3 kDa (166 aa, 1-146 aa + His Taq).

**STORAGE CONDITIONS:** Can be stored at 4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.


**DESCRIPTION**: CDA (Cytidine deaminase) is an enzyme that scavenges exogenous and endogenous cytidine and 2'-deoxycytidine for UMP synthesis. This protein is one of several deaminases responsible for maintaining the cellular pyrimidine pool. The protein also catalyzes the deamination of chemotherapeutic cytosine nucleoside analogs such as Ara-C and 5-azacytidine, which results in the loss of their cytotoxic and antitumor function. It can form homotetramers and is mainly expressed in granulocytes. Recombinant human CDA protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography.

AMINO ACID SEQUENCE: MGSSHHHHHH SSGLVPRGSH MAQKRPACTL **KPECVQQLLV CSQEAKQSAY CPYSHFPVGA** ALLTQEGRIF **KGCNIENACY PLGICAERTA** IQKAVSEGYK DFRAIAIASD MQDDFISPCG **ACROVMRFFG** TNWPVYMTKP DGTYIVMTVQ ELLPSSFGPE DLQKTQ

**BIOLOGICAL ACTIVITY:** Specific activity is > 3.5 unit/mg. One unit of activity was defined as the amount required to deaminate 1.0 µmole of cytidine per min at pH 7.5 at  $25^{\circ}$ C

## **ACTIVITY ASSAY:**

- 1. Prepare 150  $\mu$ l assay buffer. The final concentrations are 50 mM Tris-HCl (pH 7.5), 100  $\mu$ M cytidine.
- 2. Add 50 µl recombinant CDA protein with concentration 0.25 µg in assay buffer.
- 3. Mix by inversion and records the decrease at A280 nm for approximately 5 minutes.



## Cytidine millimolar extinction coefficient:

- 5.27 (mM<sup>-1</sup>.cm<sup>-1</sup>)
- 100 µM / 50 mM Tris-HCl (pH 7.5)
- PerkinElmer VICTOR X3 / A280nm
- UV Plate (Costar, Cat. No. 3635)
- Cytidine (Sigma, Cat. No. C4654)

15% SDS-PAGE (3ug)

CDA, human recombinant

## RELATED PRODUCTS:

- 2'-Deoxy-5-fluorocytidine (Cat. No. 2387-50, -250)
- 5-Aza-2'-deoxycytidine (Cat. No. 1754-10, -50)
- 5-Azacytidine (Cat. No. 1854-50, -250)
- 5-Methylcytidine (Cat. No. 2306-50, -250)
- Clofarabine (Cat. No. 2236-5, -25)
- Gemcitabine Hydrochloride (Cat. No. 1759-25, -100)
- Tetrahydrouridine (Cat. No. 1671-10, -25)
- EZSolution™ Tetrahydrouridine (Cat. No. 2355-25)
- Zebularine (Cat. No. 2225-1, -5)

FOR RESEARCH USE ONLY! Not to be used in humans.

