## **BioVision**

FORM:

## Human CellExp<sup>™</sup> CD74/DHLAG, human recombinant

CATALOG #:	7387-10	10 µg	
	7387-50	50 µg	
ALTERNATE NAMES:	CD74, DHLAG, HLADG, II, Ia-GAMMA, p33,		
	CLIP, INVG34	LIP, INVG34	
SOURCE:	HEK 293 cells (GIn	73 –Met 232)	

PURITY: ≥ 95% by SDS-PAGE get

**MOL. WEIGHT:** This protein is fused with 6×His tag at the N-terminus, has a calculated MW of 19.3 kDa. The predicted N-terminus is Gln 73. DTT-reduced Protein migrates as 23-33 kDa due to glycosylation.

ENDOTOXIN LEVEL:	<1 EU/µg by LAL method

**FORMULATION:** Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Generally 5-8% Mannitol or trehalose is added as a protectant before lyophilization.


Lyophilized

**STORAGE CONDITIONS:** Store at -20°C. After reconstitution, aliquot and store at -20°C and use within 3 months. Avoid repeated freezing and thawing cycles.

**RECONSTITUTION:** Centrifuge the vial prior to opening. Reconstitute in sterile PBS, pH 7.4 to a concentration of 50  $\mu$ g/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 month. For extended storage, it is recommended to store at -20°C.

**DESCRIPTION**: Cluster of Differentiation 74 (CD74) also known as HLA class II histocompatibility antigen gamma chain (HLADG), is a polypeptide involved in the formation and transport of MHC class II protein. The cell surface form of the invariant chain is known as CD74. CD74 is upregulated in several cancers and is also expressed by non-immune cells during inflammation. CD74 plays a critical role in MHC class II antigen processing by stabilizing peptide-free class II alpha / beta heterodimers in a complex soon after their synthesis and directing transport of the complex from the endoplasmic reticulum to the endosomal / lysosomal system where the antigen processing and binding of antigenic peptides to MHC class II takes place. CD74 / DHLAG

**BIOLOGICAL ACTIVITY:** Measured by its binding ability in a functional ELISA. Immobilized human CD74 at 5  $\mu$ g/ml (100  $\mu$ l/well) can bind biotinylated human CTSL1 with a linear range of 3.2 - 500 ng/ml.



Human recombinant CD74/DHLAG

## **RELATED PRODUCTS:**

- Human CellExp™ CD223, human recombinant (Cat. No. 7278-10, -50)
- Human CellExp<sup>™</sup> CD71, human recombinant (Cat. No. 7279-10, -50)
- Human CellExp<sup>™</sup> CD273, human recombinant (Cat. No. 7369-10, -50)
- Human CellExp<sup>™</sup> CD33, human recombinant (Cat. No. 7370-10, -50)
- Human CellExp<sup>™</sup> CD36, human recombinant (Cat. No. 7371-10, -50)
- Human CellExp<sup>™</sup> CD87, human recombinant (Cat. No. 7372-20, -100)
- Human CellExp<sup>™</sup> CD360, human recombinant (Cat. No. 7373-20, -100)
- Human CellExp<sup>™</sup> CD244, human recombinant (Cat. No. 7374-10, -50)
- Human CellExp<sup>™</sup> CD304, human recombinant (Cat. No. 7375-10)
- Human CellExp<sup>™</sup> CD319, human recombinant (Cat. No. 7376-10, -50)
- Human CellExp<sup>™</sup> CD306, human recombinant (Cat. No. 7377-10, -50)
- Human CellExp<sup>™</sup> CD84, human recombinant (Cat. No. 7378-10, -50)
- Human CellExp<sup>™</sup> CD111, human recombinant (Cat. No. 7379-10, -50)
- Human CellExp<sup>™</sup> CD48, human recombinant (Cat. No. 7380-10, -50)

FOR RESEARCH USE ONLY! Not to be used in humans.

