

Fibrinogen (Pg, Fn & vWF depleted), Human Plasma

CATALOG #:	7694-250	250 mg
ALTERNATE NAMES:	Factor 1	
SOURCE:	Human Plasma. The Human Fibrinogen, Plasminogen, Fibronectin and vWF Depleted is homogeneous as judged on a 4-20% gradient gel and is greater than 95% clottable. The plasminogen, Fibronectin and von Willebrand Factor were depleted using affinity chromatography. Purified from Human plasma that was tested and found negative by FDA accepted methods for Anti-HIV1/2, Anti-HTLV I & II, HBsAg, Anti-HCV, Syphilis, HBC Ab, HIV-1 p24 Ag or HIV-1 RNA, HCV RNA and HBV RNA. Donors are screened for CJD (Creutzfeldt-Jakob Disease).	
MOL. WEIGHT:	330.00 kDa	
PURITY:	≥ 95% by SDS-PAGE.	
EXTINCTION COEFFICIENT (1%):	15.1	
FORM:	Liquid	
FORMULATION:	In 20 mM Sodium Citrate-HCl pH 7.4.	
ACTIVITY:	> 95% clottable by functional assays.	
STORAGE CONDITIONS:	Store at -60°C or lower. Avoid repeated freezing and thawing cycles.	

DESCRIPTION: Fibrinogen is an acute phase protein that is part of the coagulation cascade of proteins. The end result of the cascade is the production of thrombin that converts fibrinogen to fibrin. Thrombin rapidly proteolysis fibrinogen, releasing fibrinopeptide A. The loss of this small peptide is not sufficient to make the resulting fibrin molecule insoluble, but it tends to form complexes with adjacent fibrin and fibrinogen molecules. Thrombin then cleaves a second peptide, fibrinopeptide B, from fibrin and the fibrin monomers formed then polymerize spontaneously to form an insoluble gel. The polymerized fibrin is held together by

noncovalent and electrostatic forces and stabilized by the transamidating enzyme, factor XIIIa that is produced by the action of thrombin on factor XIII. The insoluble fibrin aggregates (clots) and aggregated platelets then block the damaged blood vessel and prevent further bleeding. The amount of fibrinogen in the plasma can serve as a nonspecific indicator of whether or not an inflammatory process is present in the body. Fibrinogen from any mammalian source will be cleaved by thrombin from any mammalian source.

The protein was purified from Human plasma units that test negative for antibodies to HBc, HCV, HIV 1/2, HTLV-I/II, and non-reactive for HBsAg, HIV-1 RNA, HCV RNA, HBV DNA and Syphilis by FDA approved tests. Donors are screened for CJD.

Fibrinogen MUST be thawed in a 37°C water bath. Once thawed, do not chill. Remaining protein may be aliquoted into single use size and frozen at < -60°C.

RELATED PRODUCTS:

- Fibrinogen (Plasminogen depleted), Human Plasma (**Cat. No. 7692-1**)
- Fibrinogen (Pg & vWF depleted), Human Plasma (**Cat. No. 7693-500**)

FOR RESEARCH USE ONLY! Not to be used in humans.

