

Active RAF1(EE)

Recombinant protein expressed in Sf9 cells

Catalog # 7726-5, -100

Product Description

Recombinant human RAF1 (Y340E Y341E, 306-end) was expressed by baculovirus in Sf9 insect cells using an N-terminal GST tag. The gene accession number is [NM_002880](#).

Gene Aliases

None

Formulation

Recombinant protein stored in 50mM Tris-HCl, pH 7.5, 150mM NaCl, 0.25mM DTT, 0.1mM EGTA, 0.1mM EDTA, 0.1mM PMSF, 25% glycerol.


Storage and Stability

Store product at -70°C . For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

Scientific Background


RAF1 is a MAP kinase kinase kinase (MAP3K), which functions downstream of the Ras family of membrane associated GTPases to which it binds directly (1). The activated RAF1 can phosphorylate and activate the dual specificity protein kinases MEK1 and MEK2, which in turn phosphorylate to activate the serine/threonine specific protein kinases ERK1 and ERK2. Activated ERKs are pleiotropic effectors of cell physiology and play an important role in the control of gene expression involved in the cell division cycle, apoptosis, cell differentiation

Specific Activity



The specific activity of RAF1(EE) was determined to be **-6,000 nmol/min/mg** in a coupled assay as per activity assay protocol.

Purity



The purity was determined to be **>85%** by densitometry. Approx. MW **63kDa**.

RAF1(EE), Active

Recombinant protein expressed in Sf9 cells

| | |
|---------------------|-------------------------------------|
| Catalog Number | 7726-5, -100 |
| Quantity | 5 μg ; 100 μg |
| Specific Activity | ~6,000 nmol/min/mg |
| Specific Lot Number | G190-1 |



1. Kapp, U. et al: Structure and biological activity of v-rat, a unique oncogene transduced by a retrovirus. *Proc. Nat. Acad. Sci.* 80: 4218-4222, 1983.
2. Li, P. et al: Raf-1: a kinase currently without a cause but not lacking in effects. *Cell* 64: 479-482, 1991.

Storage & Shipping Store product at -70°C . For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles. Product shipped on dry ice.

To place your order, please contact us by phone 1-(650)-428-0236, fax 1-650-428-0336 or by email: orders@biovision.com
www.biovision.com

FOR IN VITRO RESEARCH PURPOSES ONLY. NOT INTENDED FOR USE IN HUMANS OR ANIMALS.

Activity Assay Protocol

Reaction Components

Active Kinase (Catalog #: 7726-5)

Active RAF1(EE) (0.1µg/µl) diluted with Kinase Dilution Buffer III and assayed as outlined in sample activity plot. (Note: these are suggested working dilutions and it is recommended that the researcher perform a serial dilution of Active RAF1(EE) for optimal results).

Kinase Dilution Buffer III

Kinase Assay Buffer I diluted at a 1:4 ratio (5X dilution) with 50ng/µl BSA solution.

Kinase Assay Buffer I

Buffer components: 25mM MOPS, pH 7.2, 12.5mM β-glycerol-phosphate, 25mM MgCl₂, 5mM EGTA, 2mM EDTA. Add 0.25mM DTT to Kinase Assay Buffer prior to use.

[³²P]-ATP Assay Cocktail

Prepare 250µM [³²P]-ATP Assay Cocktail in a designated radioactive working area by adding the following components: 150µl of 10mM ATP Stock Solution, 100µl [³²P]-ATP (1mCi/100µl), 5.75ml of Kinase Assay Buffer I. Store 1ml aliquots at -20°C.

10mM ATP Stock Solution

Prepare ATP stock solution by dissolving 55mg of ATP in 10ml of Kinase Assay Buffer I. Store 200µl aliquots at -20°C.

Substrate

Unactive MEK1 and ERK1 were activated in a coupled reaction. Myelin Basic Protein (MBP) diluted in distilled H₂O to a final concentration of 1mg/ml was subsequently used as a substrate for the activated ERK1.

Assay Protocol

Step 1. Thaw the Active RAF1(EE), Kinase Assay Buffer, Unactive ERK1 and Unactive MEK1 on ice. In a pre-cooled microfuge tube, add the following reaction components bringing the initial reaction volume up to 20µl:

- Component 1.** 10µl of diluted Active RAF1(EE)
- Component 2.** 2µl of Unactive MEK1 (0.2µg/µl)
- Component 3.** 3µl of Unactive ERK1 (0.2µg/µl)
- Component 4.** 5µl of Kinase Dilution Buffer

Step 2. Start the reaction by the addition of 5 µl ATP (250µM) and incubate in a water bath at 30° C for 25 minutes.

Step 3. After the 25 minute incubation period, remove 5µl and add to the following reaction components bringing the initial reaction volume up to 20µl on ice:

- Component 1.** 5µl of reaction mixture
- Component 2.** 10µl distilled H₂O on ice
- Component 3.** 5µl of MBP substrate on ice (1 mg/ml)

Step 4. Set up the blank control as outlined in step 3, excluding the addition of the substrate. Replace the substrate with an equal volume of distilled H₂O.

Step 5. Initiate the reaction by the addition of 5µl [³²P]-ATP Assay Cocktail bringing the final volume up to 25µl and incubate the mixture in a water bath at 30°C for 15 minutes.

Step 6. After the 15 minute incubation period, terminate the reaction by spotting 20µl of the reaction mixture onto individual pre-cut strips of phosphocellulose P81 paper.

Step 7. Air dry the pre-cut P81 strip and sequentially wash in a 1% phosphoric acid solution (dilute 10ml of phosphoric acid and make a 1L solution with distilled H₂O) with constant gentle stirring. It is recommended that the strips be washed a total of 3 intervals for approximately 10 minutes each.

Step 8. Count the radioactivity on the P81 paper in the presence of scintillation fluid in a scintillation counter.

Gentaur Europe BVBA Voorstraat 49, 1910 Kampenhout BELGIUM
Tel 0032 16 58 90 45 info@gentaur.com



Specific activity (SA) = cpm for 5µl [³²P]-ATP / pmoles of ATP (in 5µl of a 250µM ATP stock solution, i.e., 1250 pmoles)

Kinase Specific Activity (SA) (pmol/min/µg or nmol/min/mg)

Corrected cpm from reaction / [(SA of ³²P-ATP in cpm/pmol)*(Reaction time in min)*(Enzyme amount in µg or mg)]*((Reaction Volume) / (Spot Volume)]

To place your order, please contact us by phone 1-(650)-428-0236, fax 1-650-428-0336 or by email: orders@biovision.com.com
www.biovision.com

FOR IN VITRO RESEARCH PURPOSES ONLY. NOT INTENDED FOR USE IN HUMANS OR ANIMALS.