

Product Specification

CAMK4, active

(Full-length recombinant protein expressed in Sf 9 cells)

Catalog #: 7740
Lot #: _____
Aliquot size: 5 µg protein in 50 µl
Specific activity: 83 nmol/min/mg


Quality Control Analysis

Activity assessment

CAMK4 protein (~100 ng/µl concentration) was diluted to 20ng/µl in assay dilution buffer (4 mM MOPS, pH 7.2, 2.5 mM β-glycerophosphate, 1 mM EGTA, 0.4 mM EDTA, 30 mM MgCl₂, 0.05 mM DTT and 40ng/µl BSA), followed by 2-fold serial dilutions, and then the 10µl diluted proteins were used to phosphorylate the Autocamtide 2 (KKALRRQETVDAL-amide) in the following assay condition:

- 10 µl diluted CAMK4 protein
- 7.5 µl Autocamtide 2 (1mg/ml stock)
- 2.5 µl Calmodulin (0.3 mg/ml in 5mM CaCl₂)
- 5 µl [³²P] ATP mixture (250 µM ATP stock, 0.16 µCi/µl in 4x assay dilution buffer)

The various reaction components, except [³²P] ATP, were incubated at 30° C and the reaction started by the addition of [³²P] ATP. After 15 minutes, the reaction was terminated by spotting 20 µl of the reaction mixture onto a phosphocellulose P81 paper. The P81 paper was dried and washed several times in 1% phosphoric acid prior to counting in the presence of scintillation fluid in a scintillation counter. The actual counts, using various dilutions of the enzyme in the assay, are shown in Fig. 1.



Gentaur Europe BVBA Voorstraat 49, 1910 Kampenhout BELGIUM
 Tel 0032 16 58 90 45 info@gentaur.com




Fig. 1 CAMK4 activity assay

Fig. 2 CAMK4 protein gel

Purity assessment

1 µg of CAMK4 protein was subjected to SDS-PAGE and Coomassie blue staining. The scan of the gel showed >90% purity of the CAMK4 band product, and the band was at ~79 kDa (Fig. 2).



Product Description

Recombinant full-length human CAMK4 containing N-terminal GST tag was expressed by baculovirus in Sf 9 insect cells.

The gene accession number is NM_001744.

This material is sold for research purposes only.

Specific Activity

83 nmol phosphate incorporated into Autocamtide 2 per minute per mg protein at 30° C for 15 minutes using a final concentration of 50 µM ATP (0.83 µCi/assay).

Formulation

Recombinant protein in storage buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.25 mM DTT, 0.1 mM EGTA, 0.1 mM EDTA, 0.1 mM PMSF, 25% glycerol).

Storage and Stability

Store product frozen at or below -70° C. Stable for 1 year at -70° C as undiluted stock. Aliquot to avoid repeated thawing and freezing.

Scientific Background

CAMK4 belongs to the serine/threonine protein kinase family and Ca(2+)/calmodulin-dependent protein kinase subfamily. This enzyme is a multifunctional serine/threonine protein kinase with limited tissue distribution. Sikela et al. identified that brain CaM kinase IV (CAMK4) appeared to be restricted to brain in the rat; interestingly, it was not detected in the brain of the newborn, but became detectable within a few days after birth. They also raised the possibility that a defect in Camk4 may be responsible for 1 of the mutant phenotypes in the centromeric region near 2 mutations, which known to affect neurologic function and fertility (1). Wei et al. found Camk4-labeled neurons in the hippocampus, amygdala, anterior cingulate cortex, somatosensory cortex, and insular cortex (2). Kang et al. generated transgenic mice in which the expression of a dominant-negative form of Camk4 was restricted to the postnatal forebrain. In these transgenic mice, activity-induced Creb phosphorylation and Fos expression were significantly attenuated. The results indicated that neural activity-dependent CAMK4 signaling in the neuronal nucleus plays an important role in the consolidation/retention of hippocampus-dependent long-term memory (3). Wu et al. generated transgenic mice that selectively express in skeletal muscle a constitutively active form of CAMK4 and concluded that a calcium-regulated signaling pathway controls mitochondrial biogenesis in mammalian cells (4). Wei et al. studied pain and fear memory in Camk4-null mice and concluded that Camk4 is crucial in the trapping of Ca2(+)/calmodulin complexes in neuronal nuclei and in Creb phosphorylation and activation (5).

References

1. Sikela, J. M.; Law, M. L.; Kao, F.-T.; Hartz, J. A.; Wei, Q.; Hahn, W. E.: Chromosomal localization of the human gene for brain Ca(2+)/calmodulin-dependent protein kinase type IV. *Genomics* 4: 21-27, 1989.
2. Wei, F.; Qiu, C.S.; Liauw, J.; Robinson, D. A.; Ho, N.; Chatila, T.; Zhuo, M.: Calcium-calmodulin-dependent protein kinase IV is required for fear memory. *Nature Neurosci.* 5: 573-579, 2002.
3. Kang, H.; Sun, L. D.; Atkins, C. M.; Soderling, T. R.; Wilson, M. A.; Tonegawa, S.: An important role of neural activity-dependent CaMKIV signaling in the consolidation of long-term memory. *Cell* 106: 771-783, 2001.

Gentaur Europe BVBA Voorstraat 49, 1910 Kampenhout BELGIUM
Tel 0032 16 58 90 45 info@gentaur.com



dependent protein kinase IV is required for fear memory. *Nature Neurosci.* 5: 573-579, 2002.