

Recombinant Human Liver Glucokinase

CATALOG #:	7776-10
SOURCE:	<i>E. coli</i>
PURITY:	≥77% by SDS-PAGE
MOL. WEIGHT:	53.4 kDa
FORMULATION:	Liquid in 50 mM potassium phosphate pH 7.4, 50 mM sodium chloride, 0.5 mM DTT, 0.5 mM EDTA, and 2.5% glycerol.
CONCENTRATION:	14.8 mg/ml


STORAGE CONDITIONS:

The protein is best stored in working aliquots at -80 °C. Avoid freeze/thaw cycles.

DESCRIPTION:

Glucose is phosphorylated to glucose-6-phosphate by glucokinases. This gene is alternatively spliced to generate three different forms of the enzyme; one found in the pancreas and two found in the liver. The main function of this gene is to regulate carbohydrate metabolism. Recombinant human liver Glucokinase (Accession# NP_277042) has a C-terminal FLAG tag and has 474 amino acid residues. It can be useful for studies including enzyme kinetics, activator screening and kinase selectivity.

SDS-PAGE:



ACTIVATION: (\pm) Ro 28-0450 (100 μ M) activated human liver glucokinase 868% versus control measured at glucose [S]_{0.5}.

RELATED PRODUCTS:

- **Cat# 3153-100** GCK (Glucokinase) Antibody
- **Cat# 3154-100** GCK (Glucokinase) Antibody
- **Cat# 7777-10** Recombinant Human Pancreatic Glucokinase
- **Cat# K616-100** Glucose and Sucrose Assay Kit
- **Cat# K606-100** Glucose Assay Kit
- **Cat# K686-100** Glucose Assay Kit II
- **Cat# K786-100** Glucose Dehydrogenase Activity Assay Kit
- **Cat# K657-100** Glucose-6-Phosphate Assay Kit
- **Cat# K757-100** Glucose-6-Phosphate Dehydrogenase Activity Assay Kit
- **Cat# K618-100** Maltose and Glucose Assay Kit
- **Cat# K687-100** PicoProbe™ Glucose-6-Phosphate Assay Kit

SPECIFIC ACTIVITY: 662 pmol/min/ μ g. One unit is defined as the amount of enzyme that will convert 1 pmol of NADP to NADPH at 30°C. Assay conditions: 25 mM HEPES, pH 7.5, 2 mM MgCl₂, 1.0 mM DTT, 0.5 mM NADP, 2 mM ATP, 25 mM glucose, 100 μ g/ml BSA, 20 units/ml glucose 6-phosphate dehydrogenase, and 10 nM human liver glucokinase at 30°C for 30 min.

