


Human CellExp™ Testican-1, human recombinant

CATALOG NO:	P1143-10 10 µg P1143-50 50 µg
ALTERNATE NAMES:	SPOCK, TIC1, TICN1, Testican-1
SOURCE:	HEK 293 cells (Arg 22 – Trp 439)
PURITY:	> 85% by SDS – PAGE
MOL. WEIGHT:	This protein carries a polyhistidine tag at the N-terminus. The protein has a calculated MW of 47.8 kDa. The protein migrates as 50-60 kDa on a SDS-PAGE gel under reducing (R) condition due to glycosylation
ENDOTOXIN LEVEL:	< 1.0 EU per 1µg of protein (determined by LAL method)
FORM:	Lyophilized
FORMULATION:	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Generally Mannitol or Trehalose is added as a protectant before lyophilization.
STORAGE CONDITIONS:	Store at -20°C. After reconstitution, aliquot and store at -20°C and use within 3 months. Avoid repeated freezing and thawing cycles.
RECONSTITUTION:	Centrifuge the vial prior to opening. Reconstitute in sterile deionized water to a concentration of 50 µg/ml. Solubilize for 30 to 60 minutes at room temperature with occasional gentle mixing. Carrier protein (0.1% (W/V) HSA or BSA) is recommended for further dilution and long term storage. Do not vortex. This solution can be stored at 2-8°C for up to 1 month. For extended storage, it is recommended to store at -80°C.
DESCRIPTION:	Testican-1 is also known as protein SPOCK, SPOCK1, which contains one azal-like domain and one thyroglobulin type-1 domain. SPOCK1 is an extracellular heparan/chondroitin sulfate proteoglycan. Members of this family are known as testicans, also called SPOCKs. SPOCKs are enriched in brain and have been shown to regulate neuronal attachment and outgrowth. They contain inhibitory regions in several domains targeted to different classes of protease, and in some cases may act as protease inhibitors. Little is known about SPOCK1's function until now. It may play a role in cell-cell and cell-matrix interactions except for contributing to various neuronal mechanisms in the central nervous system.



Human Testican 1, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue.

RELATED PRODUCT:

- Human CellExp™ TNFRSF10B /TRAILR2, human recombinant (Cat. No. 7448-10)
- Human CellExp™ CD155, human recombinant (Cat. No. 7462-10, -50)
- Human CellExp™ CD160/BY55, human recombinant (Cat. No. 7386-10, -50)
- Human CellExp™ CD166/ ALCAM, human recombinant (Cat. No. 7437-10, -50)
- Human CellExp™ CD172A / SIRP, human recombinant (Cat. No. 7506-10, -50)
- Human CellExp™ CD33 / SIGLEC-3, human recombinant (Cat. No. 7370-10, -50)
- Human CellExp™ CD47, human recombinant (Cat. No. 7385-10, -50)
- Human CellExp™ CD55/DAF, human recombinant (Cat. No. 7432-10, -50)
- Human CellExp™ CD58 /LFA-3, human recombinant (Cat. No. 7427-10, -50)
- Human CellExp™ CD62E/E-Selectin, human recombinant (Cat. No. 7434-20, -100)
- Human CellExp™ CD71 / TFRC / TFR, human recombinant (Cat. No. 7279-10, -50)
- Human CellExp™ CD273, human recombinant (Cat. No. 7369-10, -50)
- Human CellExp™ CD36, human recombinant (Cat. No. 7371-10, -50)
- Human CellExp™ CD87, human recombinant (Cat. No. 7372-20, -100)

FOR RESEARCH USE ONLY! Not to be used on humans.

