BioVision 04/17 For research use only

Human CellExp™ LILRB4 / CD85k / ILT3, Mouse Recombinant

CATALOG NO: P1188-10 10 μg P1188-50 50 μg

ALTERNATE NAMES: LILRB4. ILT3. LIR5. CD85K. HM18

SOURCE: HEK 293 cells (Gly 24 - Lys 238)

PURITY: > 95% by SDS – PAGE

MOL. WEIGHT: This protein carries a human IgG1 Fc tag at the C-terminus. The

protein has a calculated MW of 50.7 kDa. As a result of glycosylation, the protein migrates as 60-66 kDa under reducing (R) condition, and 120-140 kDa under non-reducing (NR) condition

(SDS-PAGE).

ENDOTOXIN LEVEL: < 1.0 EU per 1µg of protein (determined by LAL method)

FORM: Lyophilized

FORMULATION: Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Generally

Mannitol or Trehalose is added as a protectant before

lyophilization.

STORAGE CONDITIONS: Store at -20°C. After reconstitution, aliquot and store at -20°C and

use within 3 months. Avoid repeated freezing and thawing cycles.

RECONSTITUTION: Centrifuge the vial prior to opening. Reconstitute in sterile

deionized water to a concentration of 50 µg/ml. Solubilize for 30 to 60 minutes at room temperature with occasional gentle mixing. Carrier protein (0.1% (W/V) HSA or BSA) is recommended for further dilution and long term storage. Do not vortex. This solution


can be stored at 2-8°C for up to 1 month.

mobilization of intracellular calcium ions.

DESCRIPTION: Leukocyte immunoglobulin-like receptor subfamily B member 4

(LILRB4) is also known as CD85 antigen-like family member K (CD85K), Immunoglobulin-like transcript 3 (ILT-3), Leukocyte immunoglobulin-like receptor 5 (LIR-5), Monocyte inhibitory receptor HM18, which belongs to the leukocyte immunoglobulin-like receptor (LIR) family. LILRB4 / CD85K contains 2 Ig-like C2-type (immunoglobulin-like) domains. CD85K is detected in monocytes, macrophages, dendritic cells, lung, natural killer cells and B-cells. LILRB4 / CD85K is receptor for class I MHC antigens. CD85K recognizes a broad spectrum of HLA-A, HLA-B, HLA-C and HLA-G alleles, involved in the down-regulation of the immune response and the development of tolerance. LILRB4 interferes with TNFRSF5-signaling and NF-kappa-B up-regulation and inhibits receptor-mediated phosphorylation of cellular proteins and

A B






Fig. A. Mouse LILRB4, Fc Tag on SDS-PAGE under reducing (R) and noreducing (NR) conditions.

Fig. B. Immobilized Human ANGPTL7, His Tag at 5 μg/mL (100 μL/well) can bind Mouse LILRB4, Fc Tag with a linear range of 0.156-1.25 μg/mL.

SPECIFIC ACTIVITY: Immobilized Human ANGPTL7, His Tag at 5 μ g/mL (100 μ L/well) can bind Mouse LILRB4, Fc Tag with a linear range of 0.156-1.25 μ g/mL.

RELATED PRODUCT:

- Human CellExp[™] TNFRSF10B /TRAILR2, human recombinant (Cat. No. 7448-10)
- Human CellExp™ CD155, human recombinant (Cat. No. 7462-10, -50)
- Human CellExp™ CD160/BY55, human recombinant (Cat. No. 7386-10, -50)
- Human CellExp™ CD166/ ALCAM, human recombinant (Cat. No. 7437-10, -50)
- Human CellExp™ CD172A / SIRP, human recombinant (Cat. No. 7506-10, -50)
- Human CellExp™ CD33 / SIGLEC-3, human recombinant (Cat. No. 7370-10, -50)
- Human CellExp[™] CD47, human recombinant (Cat. No. 7385-10, -50)
- Human CellExp™ CD55/DAF, human recombinant (Cat. No. 7432-10, -50)
- Human CellExp™ CD58 /LFA-3, human recombinant (Cat. No. 7427-10, -50)
- Human CellExp™ CD62E/E-Selectin, human recombinant (Cat. No. 7434-20, -100)
- Human CellExp™ CD71 / TFRC / TFR, human recombinant (Cat. No. 7279-10, -50
- Human CellExp™ CD273, human recombinant (Cat. No. 7369-10, -50)
- Human CellExp™ CD36, human recombinant (Cat. No. 7371-10, -50)
- Human CellExp™ CD87, human recombinant (Cat. No. 7372-20, -100)

FOR RESEARCH USE ONLY! Not to be used on humans.

